

# REGAL FERTILIZER w/ Regal RonStar 1%

OXADIAZON

GROUP

14

HERBICIDE

## RESTRICTED USE PESTICIDE

Due to potential human health risks of concern (cancer) from chronic dietary or dermal exposure when used in accordance with label directions. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

NOT FOR USE ON HOME LAWNS. FOR SALE AND USE BY PROFESSIONAL APPLICATORS ONLY. NOT FOR SALE TO, OR USE BY, HOMEOWNERS/CONSUMERS.

PREEMERGENT HERBICIDE FOR TURF AND ORNAMENTALS Selective, preemergent herbicide for the control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines and trees.

### ACTIVE INGREDIENTS

Oxadiazon: 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-delta-2,1,3,4-oxadiazolin-5-one	1.00%
Other Ingredients	99.00%
<b>Total:</b>	<b>100.0%</b>

# 30-0-0

## GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	30.00%
0.00% Nitrate Nitrogen	
0.00% Ammoniacal Nitrogen	
9.00% Other Water Soluble Nitrogen (and/or Urea Nitrogen)	0.00%
21.00% Water Insoluble Nitrogen	
Available Phosphorus (P <sub>2</sub> O <sub>5</sub> )	0.00%
Soluble Potassium (K <sub>2</sub> O)	0.00%
Chlorine, Not more than	0.00%
DERIVED FROM: Ureaform *this product contains 33.5% slow or controlled release Nitrogen.	

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>Call a poison control center or doctor immediately for treatment advice.</li><li>Have person sip a glass of water if able to swallow.</li><li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>Move person to fresh air.</li><li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>Call a poison control center or doctor for further treatment advice.</li></ul>

### MEDICAL HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-621-5208 for emergency medical treatment information.

For Technical Support or information regarding the use of this product, call 1-800-621-5208

EPA Reg. No. 48234-14

F968

EPA Est. No. 48234-GA-1

### PRECAUTIONARY STATEMENTS WARNING

#### Hazards to Humans and Domestic Animals

Causes skin and eye irritation. Harmful if inhaled. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### Personal Protective Equipment (PPE)

Loaders, applicators, and other handlers must wear: Coveralls worn over short-sleeved shirt and short pants, Waterproof gloves, Chemical resistant footwear plus socks, Protective eye wear, Chemical resistant headgear for overhead exposure, Long sleeved shirt and long pants  
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticides get inside. Then wash body thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This pesticide is toxic to oysters and shrimp.

This chemical has properties & characteristics associated with chemical detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of oxadiazon from runoff water and sediment.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of nontarget organisms by following label directions intended to minimize drift.

#### DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. After application, user must thoroughly irrigate the site as soon as possible on the same day of application to ensure product is washed off foliage and into the soil.

#### WEED RESISTANCE MANAGEMENT

For resistance management for Regal Ronstar Plus 1 is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to Regal Ronstar Plus 1 and other Version 31Oct2023 3 Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance- management strategies should be followed. To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Regal Ronstar Plus 1 or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development.
- Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species.
- If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- Report any incidence of local non-performance of this product against a particular weed species to your Regal Ronstar Plus 1 representative at (800) 621-5208, [and/or at <https://www.regalchem.com>].

#### BEST MANAGEMENT PRACTICES

- To the extent possible, use a diversified approach toward weed management. Whenever possible incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- Identify weeds present through scouting and history and understand their biology. The weed-control program should consider all of the weeds present.
- Difficult to control weeds may require sequential applications of herbicides with differing mechanisms of action.
- If resistance is suspected, treat weed escapes with an herbicide with a different MOA or use non-chemical methods to remove escapes.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry into treated areas that is permitted under Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is: Coveralls, Shoes plus socks, Waterproof gloves, Protective eyewear

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

#### STORAGE AND DISPOSAL

Storage: Do not contaminate water, food by storage or disposal. Do not store in or around the home. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### GENERAL INFORMATION

Regal Ronstar Plus I is a selective, preemergent herbicide for the control of many annual grasses and broadleaf weeds in turf and on a wide variety of woody ornamental Shrubs, vines and trees. Regal Ronstar Plus I controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination. Apply Regal Ronstar Plus I with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly. Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

#### RESTRICTIONS

##### For all use sites:

The minimum retreatment interval (RTI) between applications is 60 days. Total applications of oxadiazon of all physical forms must not exceed 6.0 lbs a.i./A/year if this product is used on the same site with liquid formulations.

##### For oxadiazon use on turf:

A maximum single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A. For applications on non-agricultural turf (e.g., golf courses, parks, athletic fields) that follow a liquid application, this product must be applied 60 days or later after the liquid application. After applying this product, no liquid applications of oxadiazon are allowed within the same calendar year on the same area of turf. For applications to non-agricultural turf (e.g., golf courses, parks, athletic fields), this product may not be applied prior to any liquid applications of oxadiazon within the same calendar year on the same area of turf. On golf course turf only: In areas of heavy weed infestation only, the maximum single application rate is 4.0 pounds of active ingredient per acre, and the maximum annual application rate is 8.0 pounds of active ingredient per acre. No liquid applications may be applied on areas of turf treated at this maximum application rate in the same year.

On golf course turf only: total applications of oxadiazon of all physical forms are limited to 30% of the total managed turf area of a golf course per year. Applications may be made to any turf surface type, including fairways, tees, greens, roughs, non-playable turf surfaces, or other managed turf areas. This 30% limit may be applied to the same or different turf areas from year to year. On golf course turf and sod farms: a minimum 10-foot-wide untreated vegetative buffer must be maintained between the treated area and any surface water features.

##### For ornamental landscaping, trees, shrubs, and nurseries:

A max. single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A per crop.

#### TURF

For use in established perennial Bluegrass, Bentgrass, Burmudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue and Zoysia turf; such as, fairways, parks, golf courses and lawns. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. Do not apply to wet turf. After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

# ITEM#: 5550

**PREEMRGENT WEED CONTROL**

Weeds Controlled	Amount of Regal RonStar Plus 1		Directions
	Per Acre	Per 1,000 sq. ft.	
Crabgrass (Digitaria spp.) Goosegrass (Eleusine indica) Field Sandbur (Cenchrus incertus) Annual Sedge (Cyperus compressus)	200-300 lbs.	4 1/2 – 6 3/4 lbs.	Apply late winter or early spring prior to seed germination. Application made after March or April may not be fully effective.
Annual Bluegrass (Poa Annua spp.)	300 lbs.	6 3/4 lbs.	Apply in late summer to early fall prior to seed germination.
Virginia Buttonweed (Diodia viriniana) Pigweed (Amaranthus spp.) Florida Pusley (Richardia scabra) Oxalis (Oxalis spp.) Stinging Nettle (Urtica dioica) Carpetgrass (Axonopus offinis)	300 lbs.	6 3/4 lbs.	Apply in early spring prior to seed germination.

**FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF**

Weeds Controlled	Amount of Regal RonStar Plus 1		Directions
	Per Acre	Per 1,000 sq. ft.	
Crabgrass, Goosegrass, and other annual weeds	300 lbs.	6 3/4 lbs.	For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue turf. Make application in the fall after turf has become dormant. For late, fall applications, seeding into treated areas should be delayed until 4 months after green up.

**ANNUAL BLUEGRASS CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED W/ PER. RYEGRASS ONLY**

Weeds Controlled	Amount of Regal RonStar Plus 1		Directions
	Per Acre	Per 1,000 sq. ft.	
Annual Bluegrass (Poa Annua spp.) and other annual weeds	200 lbs.	4 1/2 lbs.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. Regal Ronstar Plus I will only control the Poa annua spp. that is established from seed every year. Regal Ronstar Plus I should be applied no closer than 60 days prior to overseeding. NOTE: Do not use where Poa trivialis is present in overseed mix.

NOTE: In many areas, annual bluegrass exists as at least two biotypes or subspecies of Poa annua. The true annual biotype is Poa annua spp. annua which germinates every year from seed, while the biotype that behaves like a perennial is Poa annua spp. reptans. This perennial biotype can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

**ANNUAL WEED CONTROL DURING BERMUDAGRASS, ZOYSIAGRASS, SEASHORE PASPALUM AND TROPIC LALO (PASPALUM HIERONYMII) ESTABLISHMENT FROM SPRIGS**

Weeds Controlled	Amount of Regal RonStar Plus 1		Directions
	Per Acre	Per 1,000 sq. ft.	
Crabgrass, Goosegrass, and other annual weeds	200-300 lbs.	4 1/2 - 6 3/4 lbs.	For Bermudagrass, make application immediately prior to or immediately after sprigging. For Seashore Paspalum, make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after Regal Ronstar Plus I application may disrupt the herbicide barrier and reduce effectiveness. After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 300 pounds per acre (3 pounds active ingredient) of Regal Ronstar Plus I.  
Do not apply to Bentgrass turf mowed at less than 3/8 inch.  
Regal Ronstar Plus I may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.  
Avoid contact with flowers and shrubs except as recommended elsewhere on this label. Do not apply Regal Ronstar Plus I to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment except as noted above. Do not apply to Dichondra or Centipedegrass. Regal Ronstar Plus I should not be used on putting greens or tees because of the varying cultural practices used.  
When making applications to Tropic Lalo, applications must be made 0-12 days after sprigging. Do not use in areas where endangered Hawaiian Goose or nene (Branta sandvicensis), Hawaiian Duck (Anas wyvilliana), Version 31Oct2023  
Hawaiian Stilt (Himantopus mexicanus knudseni), and Hawaiian Moorhen (Gallinula chloropus sandvicensis) may feed.

**WOODY ORNAMENTAL SHRUBS, VINES, TREES AND CONIFER NURSERY APPLICATION**

It is impossible to test every species and variety of ornamental or nursery plants. Plant tolerance to Regal Ronstar Plus I has been found to be acceptable in the specified genera and species. Neither the manufacturer nor the seller has determined whether or not Regal Ronstar Plus I can safely be used on plants not specified on this label. The professional user should determine if Regal Ronstar Plus I can be used safely prior to use.  
Apply Regal Ronstar Plus I anytime during the year, prior to weed seed germination, for weed control. Regal Ronstar Plus I can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Regal Ronstar Plus I will improve weed control activity. Regal Ronstar Plus I should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Regal Ronstar Plus I. Do not apply to wet foliage or under conditions in which granules will collect on leaves. Do not apply in greenhouses as plant injury may result from co-distillation of Regal Ronstar Plus I active ingredient onto leaf surfaces in condensation.

**Amount To Use**

Ground Application — Apply Regal Ronstar Plus I broadcast to ornamentals at 200 to 300 pounds of product per acre (or 4 1/2 to 6 3/4 pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed on the following page. Remove existing weed growth before application.

**ANNUAL BROADLEAF WEEDS**

Bittercress	Bristly Oxtongue	Carpetweed	<i>*Use 300 pounds (3.0 lbs. of active ingredient) per acre for less susceptible weeds.</i>
Cheeseweed	Common Groundsel	Common Purslane	
*Evening Primrose	Fiddleneck	Fireweed	
*Galinsoga	Garden Spurge	Golden Ragwort	
Lambsquarters	** Liverwort	Niruri	
Pennsylvania Smartweed	Petty Spurge	**Prostrate Spurge	
Redroot Pigweed	Shepherdspurge	Sensitive Plant	
Sow Thistle	Speedwell	Spiny Amaranth	
Spotted Catsear	*Swinecress	Yellow Woodsorrel (Oxalis)	

**ANNUAL GRASSES**

Annual Bluegrass	Annual Sedge	*Barnyardgrass	<i>*Use 300 pounds (3.0 lbs. of active ingredient) per acre for less susceptible weeds.</i>
Bristly Foxtail	Crabgrass	Fall Panicum	
Green Foxtail	Goosegrass	**Jungle Ricegrass	
**Panama Paspalum	Ripgut Bromegrass	Wild Oats	

**CROP TOLERANCE**

Regal Ronstar Plus I is recommended for use on a wide variety of ornamental species and ground covers in container, field grown and landscape planting situations. Regal Ronstar Plus I may be used at the recommended rates on the following species.

**ESTABLISHED ORNAMENTALS AND GROUND COVERS**

Abutilon	Acacia	Agapanthus
Allamanda, Common	Andromeda	Aralia
Arborvitae	Ash, Green	Azalea
Baby's Breath	Bald Cypress	Bamboo, Feathery
Banyan, Weeping	Barberry	Beafsteak plant
Bearberry	Beech, European	Birch, Paper
Blanket, Flower	Blazing, Star	Boxwood, Japanese
Boxwood, Common	Bougainvillea	Brassaia, DwarfBuddleia
Bugleflower	Callistemon	Camellia, Sasanqua
Candytuft	Carissa	Carnation
Ceanothus	Cedar, Red	Cedar, Atlas
Cistus	Citrus, Ornamental*:	Grapefruit
Lemon	Lime	Mandarin
Tangelo	Tangerine	Temple
Cuban Shaddock	Valencia	Washington Navel
Cleyera	Coreopsis	Coromandel
Cotoneaster	Crabapple*	Crapemyrtle
Chrysanthemum	Cypress	Cypress, Leyland
Dahlia Daphne,	Fragrant Deutzia,	Slender
Dieffenbachia	Dogwood, Common	Dogwood, Redosier
Dogwood, Kousa	Douglas fir	Elm
Escallonia	Eucalyptus	Euryops, Gray Leaved
Euonymus, Spreading	Euonymus,	Wintercreeper
Falsetycypress, Hinoki	Falsetycypress, Japanese	Flowering Quince
Forsythia	Gardenia	Gazania, Trailing
Ginko	Gladiolus	Golden Raintree
Heath, Spring	Heather, Scotch	Heliotropium
emlock, Canada	Hibiscus	Honeysuckle
Holly	Ice Plant	Illima
Ivy, Algerian	Ivy, English	Jasmine, Orange
Jasmine, Primrose	Jasmine, Star	Jasmine, Showy
Juniper	Justica	Kwanzan Cherry
Lamb's Ear	Lantana, Trailing	Lapalapa
Larch, European	Laurel, English	Leopards - Bane
Leucothoe, Coast	Leucothoe, Drooping	Lily of the Nile
Lilac	Littleleaf Linden	Loquat*
Magnolia	Mahonia, Grape Holly	Maple
Metallic Plant	Mexican Orange	Mock Orange
Money Tree	Murraea	Myoporum
Myrtle	Nandina	Oak
Ochana	Ohai	Oleander
Orthosiphon	Osmanthus	Pachysandra
Palm, Bamboo	Palm, Fishtail	Palm, Licuala
Palm, Macarthur	Palm, Manac	Palm, Manila
Palm, Parlor	Palm, Rattan	Palm, Sealing Wax
Palm, Sugar	Paperbark Tree	Peony
Periwinkle, Common	Photinia, Frasier's	Poinciana
Pine	Pittosporum, Japanese	Plum, Natal
Podocarpus, Yew	Pomegranite*	Poplar
Potentilla	Privet	Protea species
Pyracantha	Raphiolepis	Redwood, Dawn
Rhododendron	Rose	Russian Olive
Sargent Cherry	Scotch Broom	Sedum
Serviceberry, Saskatoon*	Spruce	Statice
St. Johnswort, Golden	St. Johnswort,	Gokdecup
Strawberry Tree	Sumac, Staghorn	Sweet William
Syzygium	Trailing African Daisy	Viburnum
Wedelia	Wisteria, Chinese	Witchazel, Common
Woadwaxen, Common	Wooly Thyme	Xylosma
Yew		

**NEWLY TRANSPLANTED ORNAMENTALS AND GROUND COVERS**

Blanket Flower	Brassaia, Dwarf	Camellia	<i>*For use only on plants that will not bear edible fruit within one year of Regal Ronstar Plus I herbicide application.</i>
Carnation	Cedar, Red	Coromandel	
Dogwood, Flowering	Elm	Forsythia	
Gazania, Trailing	Hackberry	Ivy, Algerian	
Ivy, English	Hemlock, Canada	Holly	
Ice Plant	Juniper	Leopards - Bane	
Metallic Plant	Oak	Periwinkle, Common	
Periwinkle, Large	Pine	Poplar	
Privet	Protea Species	Spruce	
Trailing African Daisy	Wedelia	Wisteria	
Yew			

**FOREST NURSERY APPLICATION**

Seedlings of the following Species can be treated with Regal Ronstar Plus I herbicide when at least 4 weeks old: Pine – Loblolly, Eastern White, Slash

**SENSITIVE VARIETIES**

Under certain growing conditions, the plant varieties listed below may be sensitive to Regal Ronstar Plus I herbicide. Growers should evaluate usage on a small scale prior to large scale applications.

**PLANT SPECIES**

Plant Species	Varieties
Andromeda	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, (Rhododendron spp), Leuchtenfuer, Memoire John Hearrens, Hino-Crimson, White Grandeur, Everest, Girard Roberta Eichholz, Himalayan, Shangri-la (Cotoneaster spp)
Cotoneaster	Petite Red Imp, Seminole Bush
Crapemyrtle	Mediterranean Pink
Heath, Spring	Pendula
Hemlock, Canada	Red Heart
Hibiscus	Daphnoides, Nova Zembla, Rhododendron spp), Catawba Album
Rhododendron	Pumila
Spruce, Norway	Spruce, White
Spruce, White	Densata
Viburnum	Davidii

**NOTICE TO BUYER**

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal conditions. This warranty does not extend to use of this product contrary to label use directions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no warranties; expressed or implied.  
Ronstar® is a registered trademark of the Bayer CropScience.